Distribution of Coagulase-Negative Staphylococci and Antibiotic Resistance

Heechul Park^{1,2,§,*}, Sung-Bae Park^{1,2,§,*}, Junseong Kim^{1,2,*} and Sunghyun Kim^{1,2,†,**}

¹Department of Clinical Laboratory Science, College of Health Sciences,
Catholic University of Pusan, Busan 46252, Korea

²Clinical Trial Specialist Program for in Vitro Diagnostics, Brain Busan 21 Plus Program,
the Graduate School, Catholic University of Pusan, Busan 46252, Korea

Coagulase-negative staphylococci (CoNS) are a typical group of microorganisms, and the recent advances in laboratory technology and medicine has dramatically modified their significance in medical practice. CoNS, which were previously classified as normal bacterial flora, have recently been reported to be associated with serious infectious diseases, such as surgical wound infection or periprosthetic joint infection. Representative CoNS include *Staphylococcus epidermidis*, *S. haemolyticus*, and *S. saprophyticus*, which are known to cause serious problems in biomaterial-based and prosthetic device infections, as well as to cause simple urinary tract infections in sexually active women. Over the last decade, the clinical isolation rate of CoNS has been increasing, and antibiotic resistance has also been occurring. This review aimed to investigate the incidence of CoNS infection and to use the results as basic data for the management of CoNS, with a focus on the isolation rate and antibiotic resistance in clinical surgery.

Key Words: Coagulase-negative staphylococci (CoNS), Sepsis, Identification, Antibiotic resistance

INTRODUCTION

Microorganisms are closely associated with humans. Humans have co-evolved with the trillions of microbes inhabiting the human body and creating complex, body-habitat-specific, adaptive ecosystems that are finely attuned to the constantly changing host physiology (Lloyd-Price et al., 2016). Among the numerous microorganisms, representative bacteria are the main taxa of living things. Bacteria, are commonly considered as normal flora, as they have a symbiotic relationship with humans under normal circumstances; however, they can become dangerous pathogens

when the immune function is not in its normal state, such as during surgery and treatment (Marples et al., 1990).

Coagulase-negative staphylococci (CoNS) are a group of staph bacteria, which generally exist as normal flora, although several of these bacteria are potential pathogens under reduced immunity in humans (Al Tayyar et al., 2015; Argemi et al., 2019). CoNS bacteria have long been regarded as culture contaminants. However, their infective role as pathogens has become important in recent years (Natsis and Cohen, 2018). There are about 45 types of known CoNS, and the representative strains are *Staphylococcus epidermidis*, *S. haemolyticus*, and *S. saprophyticus* (Grace et al., 2019). CoNS strains are frequently isolated in hospital environ-

Received: May 31, 2021 / Revised: June 28, 2021 / Accepted: June 28, 2021

* Graduate student, ** Professor.

§ The first authors contributed equally to this work.

† Corresponding author: Sunghyun Kim. Department of Clinical Laboratory Science, College of Health Sciences, Catholic University of Pusan, Busan 46252, Korea.

Tel: +82-51-510-0560, Fax: +82-51-510-0568, e-mail: shkim0423@cup.ac.kr

©The Korean Society for Biomedical Laboratory Sciences. All rights reserved.

©This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Table 1. Prevalence of CoNS in hospital environments

Previous studies (publication year)	Dziri et al. (2016)	Al Tayyar et al. (2015)	Azih and Enabulele (2013)	Keim et al. (2011)	Begum and Anbumani (2011)	
Species	n (%)	n (%)	n (%)	n (%)	n (%)	Total, n (%)
<i>Staphylococcus epidermidis</i>	2 (2.4)	122 (54.7)	16 (26.7)	70 (36.6)	49 (43.4)	259 (38.7)
<i>Staphylococcus haemolyticus</i>	38 (45.8)	52 (23.4)	17 (28.3)	33 (17.3)	19 (16.8)	159 (23.7)
<i>Staphylococcus saprophyticus</i>	30 (36.1)	7 (3.1)	11 (18.3)	5 (2.6)	4 (3.5)	57 (8.5)
<i>Staphylococcus warneri</i>	2 (2.4)	4 (1.8)	ND*	35 (18.3)	7 (6.2)	48 (7.2)
<i>Staphylococcus simulans</i>	1 (1.2)	2 (0.9)	6 (10.0)	21 (11.0)	5 (4.4)	35 (5.2)
<i>Staphylococcus hominis</i>	ND*	13 (5.8)	ND*	4 (2.1)	11 (9.7)	28 (4.2)
<i>Staphylococcus capitis</i>	ND*	8 (3.6)	ND*	4 (2.1)	7 (6.2)	19 (2.8)
<i>Staphylococcus sciuri</i>	2 (2.4)	ND*	ND*	15 (7.9)	ND*	17 (2.5)
<i>Staphylococcus xylosus</i>	4 (4.8)	2 (0.9)	6 (10.0)	3 (1.6)	1 (0.9)	16 (2.4)
<i>Staphylococcus cohnii</i>	2 (2.4)	ND*	ND*	ND*	9 (8.0)	11 (1.6)
<i>Staphylococcus lugdunensis</i>	ND*	9 (4.0)	ND*	ND*	1 (0.9)	10 (1.5)
<i>Staphylococcus auricularis</i>	ND*	4 (1.8)	ND*	1 (0.5)	ND*	5 (0.7)
<i>Staphylococcus chromogenes</i>	ND*	ND*	3 (5.0)	ND*	ND*	3 (0.4)
<i>Staphylococcus pasteuri</i>	1 (1.2)	ND*	ND*	ND*	ND*	1 (0.1)
<i>Staphylococcus arlettae</i>	1 (1.2)	ND*	ND*	ND*	ND*	1 (0.1)
<i>Staphylococcus schleiferi</i>	ND*	ND*	1 (1.7)	ND*	ND*	1 (0.1)
Total, n (%)	83 (12.4)	223 (33.3)	60 (9.0)	191 (28.5)	113 (16.9)	670 (100.0)

*ND: not detected

ments; the representative CoNS strains include *S. cohnii*, *S. warneri*, *S. sciuri*, *S. hominis*, *S. simulans*, *S. pasteuri*, *S. arlettae*, and *S. xylosus* (Table 1) (Al Tayyar et al., 2015; Houssaini et al., 2019; Dziri et al., 2016; Keim et al., 2011; Begum and Anbumani, 2011). Among the CoNS species associated with clinical disease, *S. epidermidis* is identified in infection associated with biomaterial-based and prosthetic devices. Meanwhile, *S. saprophyticus* causes simple urinary tract infection in sexually active women, and *S. haemolyticus* causes variable infection such as periprosthetic joint infections (PJI) and bacteremia (Eltwisy et al., 2020). In addition, identification of CoNS is of increasing importance in infection, as the isolation rate of CoNS from the blood of patients with risk factors has been increasing (Nickel and Costerton, 1992; Houssaini et al., 2019).

1. Importance of CoNS

In recent years, the use of devices such as intravascular catheters as well as the increase in invasive manipulation in

immunosuppressed patients, have been recognized as one of the important factors associated with nosocomial infections caused by bacteria (Supré et al., 2011). The frequency of CoNS infection has been increasing mainly due to bacteremia, catheter-related infections, endocarditis, and urinary tract infections; moreover, resistance to antibiotics, especially vancomycin, is a serious problem, as bacteria demonstrating severe resistance have been identified (Koksai et al., 2009; De Vecchi et al., 2018). A 2013 study for Azih et al., has reported that 105 CoNS were isolated from 79 clinical specimens, such as discharge from wounds, urine, and male genital infections, which could cause secondary infections, such as sepsis (Azih and Enabulele, 2013). These CoNS infections are usually internal or implanted foreign bodies that cause immune damage and are emerging as the cause of infectious diseases in both the community and hospital environment (Chu et al., 2008; Piette and Verschraegen, 2009).

Table 2. Prevalence of potential pathogens isolated from the blood culture in the Republic of Korea

Microorganism	No.	True pathogen (%)	Contaminated (%)	Reference
Coagulase-negative staphylococci	560	133 (23.8)	427 (76.2)	Chun et al., 2019
Viridans group <i>Streptococcus</i>	93	43 (46.2)	50 (53.8)	
<i>Bacillus</i> species	54	15 (27.7)	39 (72.3)	
<i>Corynebacterium</i> species	21	4 (19.0)	17 (81.0)	
<i>Micrococcus</i> species	14	0 (0.0)	14 (100.0)	
<i>Propionibacterium acnes</i>	8	0 (0.0)	8 (100.0)	
Total	750	195	555	

2. Distribution of CoNS in sepsis, surgical wound infection (SWI), and periprosthetic joint infection (PJI)

2.1. CoNS in Sepsis

Sepsis is one of the most common causes of death among hospitalized patients, and it can be caused by bacteria, fungi, or viruses (Rello et al., 2017). The causative agent of sepsis varies, but 11 pathogens are known as the major causative organisms (Liu et al., 2018). *S. aureus* and *Escherichia coli* are the most common causes; CoNS is also frequently isolated from blood, which in turn is considered contaminated, although no symptoms have been reported in such cases (Biedenbach et al., 2004). However, in developed countries, CoNS, such as *S. epidermidis* and *S. haemolyticus*, are indicated as the major pathogens involved in late-onset sepsis (Golińska et al., 2020) and are among the pathogenic bacteria implicated in catheter-related blood stream infection (CRBSI) (Osaki et al., 2020). According to one report, approximately 10% of the causative pathogens of sepsis are CoNS (Chun et al., 2015), and the most detected CoNS was *S. epidermidis* (20%), followed by *S. hominis*; *S. aureus* was detected at a rate of approximately 9% (Osaki et al., 2020). In Korea, in 13,519 blood cultures obtained from pediatric patients, 750 bacteria were identified, 560 (74.67%) of which were CoNS (Table 2). Indeed, the degree of contamination is high, and this finding is significant because contaminations are highly associated with diseases (Chun et al., 2019).

2.2. CoNS in surgical wound infection (SWI)

SWI is one of the complications in variable surgery, and the risk of SWI is 1.9 per 100 interventions (Hijas-Gómez et al., 2017). Also, SWI remains as the most common and

expensive hospital-acquired infection, accounting for 20% of cases (Campwala et al., 2019). During infection, pain, swelling, erythema, warmth, and impairment of function are consistently seen (Peel and Taylor, 1991).

In one study involving SWI patients, 82 samples were analyzed; CoNS were identified in 54.9% of the samples, especially methicillin-resistant CoNS (MR-CoNS) (13.4%) (Ahmed et al., 2021). In another study, *S. aureus* (53.2%) and CoNS (23.4%) were identified in 74 SWI patients (Liang et al., 2019). In yet another study, 31 of the 52 patients were confirmed to have bacterial growth, and 20 of them were infected with *S. epidermidis* (65%). These results suggest that in SWI patients, the proportion of CoNS is quite high, and this phenomenon remains a serious healthcare-associated infection (HAI) problem.

2.3. CoNS in periprosthetic joint infection (PJI)

PJI carries both a higher economic burden and greater morbidity and mortality compared with other aseptic complications (Gross et al., 2021). A PJI infection requires extended periods of hospitalization and re-operations and thus poses a significant financial burden (Li et al., 2020). Therefore, management of established PJI requires a combination of pathogen-specific antibiotics and surgical intervention (De Vecchi et al., 2018).

These PJIs are frequently caused by CoNS; importantly, antibiotic resistance among causative pathogens of PJI has been increasing (Veltman et al., 2019). A study involving 50 CoNS and 39 *S. aureus* infection cases has found that the CoNS were significantly more resistant to daptomycin and gentamicin and were more susceptible to rifampicin than *S. aureus* (De Vecchi et al., 2018). Another study reported

that in 106 PJI cases, 43 cases involved *S. aureus* and 32 involved streptococci infections; only 8 cases involved CoNS infections (Rakow et al., 2019). Another study involving 74 patients showed that PJI was caused by methicillin-resistant *S. aureus* (MRSA) (20.3%), methicillin-resistant *S. epidermidis* (MRSE) (32.4%), methicillin-susceptible *S. aureus* (MSSA) (18.9%), and methicillin-susceptible *S. epidermidis* (MSSE) (28.4%) (Hischebeth et al., 2019). These results suggested that in PJI, the major causative agents are gram-positive bacteria, especially *S. aureus* and CoNS. Moreover, the incidence of MRSE has been increasing; thus, MRSE must be diagnosed and treated promptly.

3. Distribution of antibiotic-resistant CoNS

As mentioned previously, the distribution of CoNS in various diseases as well as the incidence of antibiotic-resistant CoNS have been increasing. In addition, biofilm formation has been observed among CoNS, making them more difficult to treat because of their antibiotic resistance compared with non-biofilm bacteria (Saber et al., 2017). Methicillin resistance in staphylococci is caused by the expression of PBP2a (PBP') encoded by the *mecA* gene (Hanssen et al., 2004). According to one study, *S. epidermidis* is the most frequent isolate among methicillin-resistant CoNS (Schuster et al., 2018). Oxacillin resistance is often mediated by the *mecA* gene, which encodes a supplemental penicillin binding protein (PBP2a) with low affinity to semisynthetic penicillins (Archer and Niemeyer, 1994). The *mecA* gene is located on a mobile genetic element known as the staphylococcal cassette chromosome *mec* (SCC*mec*) that comprises the *mec* complex consisting of the *mecA* gene and its regulator genes *mecI* and *mecRI*, along with the *ccr* complex, which is responsible for integration (Pereira et al., 2020). Among the several MR-CoNS, *S. epidermidis* and *S. haemolyticus* were the most common species identified (Venugopal et al., 2019). Also, according to one study, *S. haemolyticus* isolates demonstrated the overall highest resistance rates (Tekeli et al., 2020). Another study investigated the antibiotic resistance and virulence factors of CoNS isolated from blood cultures. Among the 93 CoNS species, 86 were resistant to cefoxitin, and 49 displayed multi-antibiotic resistance (Al-Haqan et al., 2020). Also,

vancomycin-resistant CoNS is a recent health concern, especially in serious infections, such as bloodstream infections, as it may lead to failure of therapy (Mashaly and El-Mahdy, 2017). Therefore, early detection of antibiotic-resistant CoNS is important in order to effectively eliminate the bacteria that causes bloodstream infection.

CONCLUSION

The management of the CoNS plays an important role in controlling *S. aureus*-related infectious diseases in recent years. CoNS has a symbiotic relationship with humans in a normal environment, but it can become a dangerous pathogen when immune function is not normal, such as surgery or treatment. Besides, if CoNS are identified in blood cultures, it remains a serious healthcare problem whether infection or a contaminant (Muñoz-Gamito et al., 2020). Recently, it was confirmed that the treatment for surgical wound infection and the separation rate of CoNS from the joint around the prosthesis were increased, and the infection rate was also increased in septic patients. Also, the abuse and misuse of antibiotics, and antibiotic-resistant CoNS, which is caused when appropriate treatment was not conducted, are also becoming a problem in the community. After all, CoNS has been regarded as a contaminant, however its importance has increased in relation to various clinical diseases. Therefore, a research of CoNS on the distribution in variable diseases and antibiotic-resistant patterns are important to dissolve these situations. In addition, it is necessary to characterize the role of virulence factor and pathogenesis of CoNS in the host to identify the etiology in human.

ACKNOWLEDGEMENT

This paper was supported by RESEARCH FUND from Catholic University of Pusan.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

Ahmed E, Gad G, Soliman W, El-Asady R, Hasaneen A,

- Abdelwahab SF. Prevalence of methicillin-resistant coagulase-negative staphylococci among Egyptian patients after surgical interventions. *Trop Doct*. 2021. 51: 40-44.
- Al Tayyar I, Al-Zoubi M, Hussein E, Khudairat S, Sarosiekf K. Prevalence and antimicrobial susceptibility pattern of coagulase-negative staphylococci (CoNS) isolated from clinical specimens in Northern of Jordan. *Iran J Microbiol*. 2015. 7: 294-301.
- Al-Haqan A, Boswihi SS, Pathan S, Udo EE. Antimicrobial resistance and virulence determinants in coagulase-negative staphylococci isolated mainly from preterm neonates. *Plos One*. 2020. 15: e0236713.
- Archer G, Niemeyer D. Origin and evolution of DNA associated with resistance to methicillin in staphylococci. *Trends Microbiol*. 1994. 2: 343-347.
- Argemi X, Hansmann Y, Prola K, Prevost G. Coagulase-negative staphylococci pathogenomics. *Int J Mol Sci*. 2019. 20: 1215.
- Azih A, Enabulele I. Species distribution and virulence factors of coagulase negative staphylococci isolated from clinical samples from the University of Benin Teaching hospital, Edo State, Nigeria. *J Nat Sci Res*. 2013. 3: 38-43.
- Begum E, Anbumani N. Prevalence and antimicrobial susceptibility pattern of Coagulase-negative *Staphylococcus*. *Int J Med Public Health*. 2011. 1: doi:10.5530/ijmedph.4.2011.14.
- Biedenbach D, Moet G, Jones R. Occurrence and antimicrobial resistance pattern comparisons among bloodstream infection isolates from the SENTRY Antimicrobial Surveillance Program (1997-2002). *Diagn Microbiol Infect Dis*. 2004. 50: 59-69.
- Campwala I, Unsell K, Gupta S. A comparative analysis of surgical wound infection methods: predictive values of the CDC, ASEPSIS, and Southampton scoring systems in evaluating breast reconstruction surgical site infections. *Plast Surg*. 2019. 27: 93-99.
- Chu VH, Woods CW, Miro JM, Hoen B, Cabell CH, Pappas PA, et al. Emergence of coagulase-negative staphylococci as a cause of native valve endocarditis. *Clin Infect Dis*. 2008. 46: 232-242.
- Chun K, Syndergaard C, Damas C, Trubey R, Mukindaraj A, Qian S, et al. Sepsis pathogen identification. *J Lab Autom*. 2015. 20: 539-561.
- Chun S, Kang CI, Kim YJ, Lee NY. Clinical Significance of Isolates Known to Be Blood Culture Contaminants in Pediatric Patients. *Medicina*. 2019. 55: 696.
- De Vecchi E, George DA, Romanò CL, Pregliasco FE, Mattina R, Drago L. Antibiotic sensitivities of coagulase-negative staphylococci and *Staphylococcus aureus* in hip and knee periprosthetic joint infections: does this differ if patients meet the International Consensus Meeting Criteria? *Infect Drug Resist*. 2018. 11: 539-546.
- Dziri R, Klibi N, Lozano C, Ben Said L, Bellaaj R, Tenorio C, Boudabous A, et al. High prevalence of *Staphylococcus haemolyticus* and *Staphylococcus saprophyticus* in environmental samples of a Tunisian hospital. *Diagn Microbiol Infect Dis*. 2016. 85: 136-140.
- Houssaini ZE, Harrar N, Zerouali K, Belabbes H, Elmdaghri N. Prevalence of Coagulase-Negative Staphylococci in Blood Cultures at the Ibn-Rochd University Hospital in Casablanca. *Pan Afr Med J*. 2019. 33: 193.
- Eltwisy HO, Abdel-Fattah M, Elsis AM, Omar MM, Abdelmoteleb AA, El-Mokhtar MA. Pathogenesis of *Staphylococcus haemolyticus* on primary human skin fibroblast cells. *Virulence*. 2020. 11: 1142-1157.
- Golińska E, Strus M, Tomusiak-Plebanek A, Więcek G, Kozieł Ł, Lauterbach R. Coagulase-Negative Staphylococci Contained in Gut Microbiota as a Primary Source of Sepsis in Low-and Very Low Birth Weight Neonates. *J Clin Med*. 2020. 9: 2517.
- Grace JA, Olayinka BO, Onaolapo JA, Obaro SK. *Staphylococcus aureus* and coagulase-negative staphylococci in bacteraemia: the epidemiology, predisposing factors, pathogenicity and antimicrobial resistance. *Clin Microbiol*. 2019. 8: 1-5.
- Gross CE, Della Valle CJ, Rex JC, Traven SA, Durante EC. Fungal Periprosthetic Joint Infection: A Review of Demographics and Management. *J Arthroplasty*. 2021. 36: 1758-1764.
- Hanssen AM, Kjeldsen G, Sollid JU. Local variants of Staphylococcal cassette chromosome mec in sporadic methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative Staphylococci: evidence of horizontal gene transfer?. *Antimicrob Agents Chemother*. 2004. 48: 285-296.
- Hijas-Gómez AI, Egea-Gámez RM, Martínez-Martín J, González-Díaz R, Losada-Viñas JJ, Rodríguez-Caravaca G. Surgical Wound Infection Rates and Risk Factors in Spinal Fusion in a University Teaching Hospital in Madrid, Spain. *Spine (Phila Pa 1976)*. 2017. 42: 748-754.
- Hischebeth GT, Randau TM, Ploeger MM, Friedrich MJ, Kaup E, Jacobs C, Molitor E, Hoerauf A, Gravius S, Wimmer MD. *Staphylococcus aureus* versus *Staphylococcus epidermidis* in periprosthetic joint infection-Outcome analysis of methicillin-resistant versus methicillin-susceptible strains. *Diagn Microbiol Infect Dis*. 2019. 93: 125-130.
- Keim LS, Torres-Filho SR, Silva PV, Teixeira LA. Prevalence,

- aetiology and antibiotic resistance profiles of coagulase negative staphylococci isolated in a teaching hospital. *Braz J Microbiol.* 2011. 42: 248-255.
- Koksall F, Yasar H, Samasti M. Antibiotic resistance patterns of coagulase-negative staphylococcus strains isolated from blood cultures of septicemic patients in Turkey. *Microbiol Res.* 2009. 164: 404-410.
- Li C, Renz N, Trampuz A, Ojeda-Thies C. Twenty common errors in the diagnosis and treatment of periprosthetic joint infection. *Int Orthop.* 2020. 44: 3-14.
- Liang Z, Rong K, Gu W, Yu X, Fang R, Deng Y, Lu L. Surgical site infection following elective orthopaedic surgeries in geriatric patients: Incidence and associated risk factors. *Int Wound J.* 2019. 16: 773-780.
- Liu CF, Shi XP, Chen Y, Jin Y, Zhang B. Rapid diagnosis of sepsis with TaqMan-Based multiplex real-time PCR. *J Clin Lab Anal.* 2018. 32: e22256.
- Lloyd-Price J, Abu-Ali G, Huttenhower C. The healthy human microbiome. *Genome Med.* 2016. 27. 8: 51.
- Marples RR, Richardson JF, Newton FE. Staphylococci as part of the normal flora of human skin. *Soc Appl Bacteriol Symp Ser.* 1990. 19: 93S-99S.
- Mashaly GE, El-Mahdy RH. Vancomycin heteroresistance in coagulase negative *Staphylococcus* blood stream infections from patients of intensive care units in Mansoura University Hospitals, Egypt. *Ann Clin Microbiol Antimicrob.* 2017. 19: 16: 63.
- Muñoz-Gamito G, Cuchí E, Roigé J, Gómez L, Jaén À, Pérez J, et al. Higher accuracy of genotypic identification compared to phenotyping in the diagnosis of coagulase-negative staphylococcus infection in orthopedic surgery. *Infect Dis (Lond).* 2020. 52: 883-890.
- Natsis NE, Cohen PR. Coagulase-Negative Staphylococcus Skin and Soft Tissue Infections. *Am J Clin Dermatol.* 2018. 19: 671-677.
- Nickel JC, Costerton JW. Coagulase-negative staphylococcus in chronic prostatitis. *J Urol.* 1992. 147: 398-400.
- Osaki S, Kikuchi K, Moritoki Y, Motegi C, Ohyatsu S, Murakawa Y, et al. Distinguishing coagulase-negative Staphylococcus bacteremia from contamination using blood-culture positive bottle detection pattern and time to positivity. *J Infect Chemother.* 2020. 26: 672-675.
- Peel AL, Taylor EW. Proposed definitions for the audit of post-operative infection: a discussion paper. *Surgical Infection Study Group. Ann R Coll Surg Engl.* 1991. 73: 385-388.
- Pereira VC, Romero LC, Pinheiro-Hubinger L, Oliveira A, Martins KB, Cunha MLRSD. Coagulase-negative staphylococci: a 20-year study on the antimicrobial resistance profile of blood culture isolates from a teaching hospital. *Braz J Infect Dis.* 2020. 24: 160-169.
- Piette A, Verschraegen G. Role of coagulase-negative staphylococci in human disease. *Vet Microbiol.* 2009. 16; 134: 45-54.
- Rakow A, Perka C, Trampuz A, Renz N. Origin and characteristics of haematogenous periprosthetic joint infection. *Clin Microbiol Infect.* 2019. 25: 845-850.
- Rello J, Valenzuela-Sánchez F, Ruiz-Rodriguez M, Moyano S. Sepsis: A Review of Advances in Management. *Adv Ther.* 2017. 34: 2393-2411.
- Saber H, Jasni AS, Jamaluddin TZMT, Ibrahim R. A Review of Staphylococcal Cassette Chromosome mec (SCCmec) Types in Coagulase-Negative Staphylococci (CoNS) Species. *Malays J Med Sci.* 2017. 24: 7-18.
- Schuster D, Josten M, Janssen K, Bodenstein I, Albert C, Bierbaum G, et al. Detection of methicillin-resistant coagulase-negative staphylococci harboring the class A mec complex by MALDI-TOF mass spectrometry. *Int J Med Microbiol.* 2018. 308: 522-526.
- Supré K, Haesebrouck F, Zadoks RN, Vanechoutte M, Piepers S, De Vlieghe S. Some coagulase-negative *Staphylococcus* species affect udder health more than others. *J Dairy Sci.* 2011. 94: 2329-2340.
- Tekeli A, Öcal DN, Dolapçı İ. Detection of sasX Gene and Distribution of SCCmec Types in Invasive and Non-invasive Coagulase-negative Staphylococci. *Balkan Med J.* 2020. 1; 37: 215-221.
- Veltman ES, Moojen DJF, van Ogtrop ML, Poolman RW. Two-stage revision arthroplasty for coagulase-negative staphylococcal periprosthetic joint infection of the hip and knee. *World J Orthop.* 2019. 18; 10: 348-355.
- Venugopal N, Mitra S, Tewari R, Ganaie F, Shome R, Shome BR, et al. Molecular detection and typing of methicillin-resistant *Staphylococcus aureus* and methicillin-resistant coagulase-negative staphylococci isolated from cattle, animal handlers, and their environment from Karnataka, Southern Province of India. *Vet World.* 2019. 12: 1760-1768.

<https://doi.org/10.15616/BSL.2021.27.2.45>

Cite this article as: Park H, Park SB, Kim J, Kim S. Distribution of Coagulase-Negative Staphylococci and Antibiotic Resistance. *Biomedical Science Letters.* 2021. 27: 45-50.